

**Ecography**

**ECOG-00965**

Yang, Q., Wei, S., Shang, L., Carillo, J., Gabler, C. A., Nijjer, S., Li, B. and Siemann, E. 2015. Mycorrhizal associations of an invasive tree are enhanced by both genetic and environmental mechanisms. – Ecography doi: 10.1111/ecog.00965

**Supplementary material**

## Appendix 1

Table A1. The geographical locations of *Triadica sebifera* populations from the native (China) and introduced (USA) ranges that were used in this study.

### Native populations from China

ID	Site of seed collection	Longitude	Latitude
C-AB-1	Bengbu, Anhui	32° 57' N	117° 20' E
C-FJ-1	Fuzhou, Fujian	26° 04' N	119° 17' E
C-GD-3	Zhaoqing, Guangdong	23° 03' N	112° 28' E
C-GX-1	Guilin, Guangxi	25° 15' N	110° 21' E
C-GZ-3	Dushan, Guizhou	25° 51' N	107° 31' E
C-HB-1	Wuhan, Hubei	31° 03' N	115° 25' E
C-HE-1	Sikeshuxiang, Henan	33° 18' N	112° 21' E
C-HN-2	Zhongjiapuxiang, Hunan	29° 13' N	111° 05' E
C-JN-1	Nanjing, Jiangsu	32° 05' N	118° 47' E
C-SC-1	Daweizhen, Sichuan	29° 20' N	103° 17' E
C-ST-1	Laiwu, Shandong	36° 09' N	117° 09' E
C-SX-1	Chongming Island, Shanghai	31° 38' N	121° 33' E

### Invasive populations from USA

ID	Site of seed collection	Longitude	Latitude
U-AL-1	Tillman's Corner, Alabama	30° 35' N	88° 09' W
U-FL-2	Gainseville, Florida	29° 35' N	82° 21' W
U-FL-3	Callahan, Florida	30° 35' N	81° 48' W
U-GA-1	Hutchinson Island, Georgia	32° 06' N	81° 06' W
U-GA-2	Sapelo Island, Georgia	31° 23' N	81° 15' W
U-LA-1	Lake Charles, Louisiana	30° 14' N	93° 09' W
U-LA-4	Baton Rouge, Louisiana	30° 23' N	91° 09' W
U-MS-1	Moss Point, Mississippi	30° 26' N	88° 31' W
U-SC-1	Limehouse, South Carolina	32° 09' N	81° 06' W
U-TX-2	La Marque, Texas	29° 22' N	95° 02' W
U-TX-4	Lynchburg, Texas	29° 47' N	95° 02' W
U-TX-5	Port Arthur, Texas	29° 53' N	94° 02' W