

Ecography

ECOG-01020

Blasco-Costa, I., Rouco, C. and Poulin, R. 2014.
Biogeography of parasitism in freshwater fish: spatial
patterns in hot spots of infection. – Ecography doi:
10.1111/ecog.01020

Supplementary material

Appendix 1.

Table A1. Summary data for each sampling location. Number of fish examined for parasites from each site. Favourability values for each parasite species at each site (values vary from 0 to 1).

Site	Location	No. of fish	Favourability			
			<i>Coitocaecum parvum</i>	<i>Apatemon</i> sp.	<i>Stegodexamene anguillae</i>	<i>Telogaster opisthorchis</i>
1	Taiura River	5	0.195	0.828	0.828	0.828
2	Waipapa River	10	0.107	0.783	0	0
3	Opotiki Stream	5	0.107	0.649	0.649	0.649
4	Patea River	5	0.903	0.916	0.916	0.916
5	Tuaraki River	11	0.472	0.787	0.787	0.787
6	Rangitawa Stream	10	0.759	0.759	0.759	0.759
7	Porangahau River	10	0.723	0.953	0.953	0.953
8	Silverstream Creek	10	0.54	0.79	0.79	0.79
9	Bocketts Stream	7	0.107	0.267	0.267	0.267
10	Moutere River	5	0.54	0.828	0.828	0.828
11	Rai River	9	0.195	0.849	0.849	0.849
12	Grey River	10	0.195	0.743	0.743	0.743
13	Bruce Creek	9	0.195	0.783	0.783	0.783
14	Hotikiti River	9	0.107	0.334	0.334	0.334
15	Cass Creek	6	0.107	0.294	0.294	0.294
16	Acheron River	8	0.195	0.211	0.09	0.09
17	Hawkins River	12	0.618	0.903	0.843	0.843
18	Ohau River	12	0.107	0.719	0.354	0.354
19	Manuherikia River	11	0.723	0.965	0.351	0.351
20	Aparima River	13	0.836	0.836	0.836	0.836