

## Supplementary material

### Additional species potentially occurring in TNP and CNP based on their distribution ranges, habitat preferences, and records from neighbouring sites

TNP – 4–10 species: *Nycteris nana*, *Hipposideros marisae*\*, *Kerivoula lanosa*, *Neoromicia nana*, *Neoromicia tenuipinnis*, *Glauconycteris superba*\*, *Scotophilus nucella*\*, *Myopterus whitleyi*\*, *Mops petersoni*\*, *Mops trevori*.

CNP – 8–14 species: *Taphozous perforatus*\*, *Taphozous nudiventris*\*, *Coleura afra*\*, *Nycteris grandis*, *Lavia frons*, *Rhinolophus fumigatus*\*, *Kerivoula lanosa*, *Myotis bocagii*, *Pipistrellus rusticus*, *Neoromicia capensis*\*, *Nycticeinops schlieffenii*, *Scotoecus albofuscus*\*, *Scotophilus nigrita*, *Mops midas*.

\* denotes species less likely to be found. The likelihood of local occurrence of several species partly depends on the availability of suitable day roosts, in particular caves or similar structures such as rock boulders (*Taphozous perforatus*, *T. nudiventris*, *Coleura afra*, *Rhinolophus fumigatus*, *Hipposideros marisae*).