

Ecography

E5532

Tablado, Z., Revilla, E. and Palomares, F. 2009. Breeding like rabbits: global patterns of variability and determinants of European wild rabbit reproduction. – *Ecography* 32: 310–320.

## Supplementary material

See next page.

Table S1. List of publications reviewed for this study. Last column shows the type of analyses in which their data was used.

Study	Location	Latitude	Analyses
Andersson and Meurling (1977)	Southern Sweden	55.7°N	
Andersson et al. (1981)	Southern Sweden	55.7°N	6
Arques (2000)	Eastern Spain	38.0°N	1,3,4,6
Arthur (1980)	Northern France	48.7°N	1,3,4,6
Bell (1977)	North Canterbury (New Zealand)	42.7°S	1,3
Bonino (2006)	Argentina	36.5°–55.0°S	
Brambell (1944)	Caernarvonshire (UK)	53.0°N	1,3,6
Calvete et al. (2002)	North eastern Spain	41.7°N	
Calzada (2000)	South western Spain	37.1°N	6
Delibes and Calderon (1979)	South western Spain	37.1°N	2,6
Dunsmore (1971)	Eastern coastal Australia	35.8°S	1,3,5,6
Dunsmore (1974)	Subalpine Australia	36.3°S	6
Fraser (1988)	Central Otago (New Zealand)	45.3°S	2,6
Gibb and Williams (1994)	New Zealand	35.0°–46.8°S	
Gibb et al. (1985)	Wairarapa (New Zealand)	41.2°S	1,3,6
Gonçalves et al. (2002)	Southern Portugal	38.8°N	6
Jaksic and Fuentes (1991)	Chile	29.0°–54.0°S	
Lloyd (1970)	Wales (UK)	52.5°N	
	Skokholm (UK)	51.5°N	
	Skerries (UK)	53.3°N	
Myers (1971)	Subtropical Australia	26.3°S	6
	Subalpine Australia	36.3°S	6
	Arid Australia	30.2°S	6
	Mediterranean Australia	35.3°S	6
Parer (1977)	Mediterranean Australia	35.3°S	
Parer and Libke (1991)	South eastern Australia	35.3°S	1,3,4,6
Phillips et al. (1952)	Wales (UK)	51.9°N	6
Reddiex et al. (2002)	North Canterbury (New Zealand)	42.7°S	
Ribeiro (1983)	Southern Portugal	38.1°N	1,3,4,6
Rodel et al. (2004b)	Bayreuth (Germany)	49.9°N	
Rodel et al. (2005)	Bayreuth (Germany)	49.9°N	
Rogers (1981)	Southern France	43.7°N	6
Rogers et al. (1994)	Continental Europe		
Skira (1978)	Subantarctic Australia	54.6°S	1,3,6
Soriguer (1981)	South western Spain	37.9°N	1,3,4,6
Stevens and Weisbrod (1979)	San Juan Island (USA)	48.5°N	6
Trout and Smith (1998)	Lincolnshire (UK)	53.1°N	
Twigg et al. (1998)	South western Australia	34.5°S	6
Villafuerte (1994)	South western Spain	37.1°N	2
von Holst et al. (1999)	Bayreuth (Germany)	49.9°N	
von Holst et al. (2002)	Bayreuth (Germany)	49.9°N	
Wallage-Drees (1983)	Holland	52.5°N	
Watson (1957)	Hawke's Bay (New Zealand)	39.8°S	6
Wheeler and King (1985)	Cape Naturaliste (SW Australia)	33.5°S	5,6
	Chidlow (SW Australia)	31.9°S	5,6
Williams and Robson (1985)	Wanganui district (New Zealand)	39.3°S	6
Wood (1980)	Arid Australia	29.7°S	
Zunino (1987)	Central Chile	33.2°S	1,3
Zunino and Vivar (1985)	Central Chile	33.2°S	6

1 = Breeding season GLMMs, 2 = Breeding season model validation, 3 = Pregnancy rate GLMM, 4 = Pregnancy rate GLMM with data on population change, 5 = Age effect on pregnancy rates, 6 = Litter size regressions.